

Table 1. Expanded statewide harvest estimates, average catch per trapper, average catch per successful trapper, and percent successful trappers by furbearer species for the 2015-16 season. There were n=859 licensed trappers in Mississippi in 2015-16.

Species	Harvest ^a	SE	Average Catch Per Trapper ^b	SE	Average Catch Per Successful Trapper	SE	n ^c	Percent Successful Trappers ^b	SE
Mink	35	7.15	0.04	0.01	1.46	0.12	13	2.8	0.52
Raccoon	14,456	808.69	16.83	0.94	20.72	1.12	376	81.21	1.23
Muskrat	334	61.88	0.39	0.07	3.83	0.62	47	10.15	0.95
Red Fox	403	44.32	0.47	0.05	2.44	0.22	89	19.22	1.24
Gray Fox	1,716	127.47	2.00	0.15	5.67	0.34	163	35.21	1.51
Bobcat	1,878	171.86	2.19	0.20	5.88	0.48	172	37.15	1.53
Opossum	7,687	508.55	8.94	0.66	16.31	1.14	254	54.86	1.57
Otter	1,850	166.59	2.15	0.19	6.92	0.53	144	31.10	1.46
Spotted Skunk	52	22.64	0.06	0.02	4.67	0.1.72	6	1.30	0.36
Striped Skunk	831	100.83	0.97	0.11	5.74	0.57	78	16.85	1.18
Coyote	5,115	379.55	5.96	0.44	12.36	0.82	223	48.16	1.58
Weasel	0	0.00	0.00	0.00	0.00	0.00	0	0	0.00
Nutria	1,273	278.63.	1.48	0.32	8.79	1.83	78	16.85	1.18
Beaver	9,432	661.36	10.98	0.77	20.34	1.30	250	54.00	1.57

^a n = 463

^b n = 463

^c n=sample size for average catches per successful trapper