

Bay Springs Lake 2024 REEL FACTS

Dustin Rodgers

Dustin.Rodgers@wfp.ms.gov Telephone (662) 840-5176

General Information: Covering approximately 6,700 acres and 8 miles in length, Bay Springs Lake is the northern most impoundment along the Tennessee-Tombigbee Waterway. Bay Springs is connected to Pickwick Lake on the Tennessee River through the Divide Cut and Yellow Creek. As a result, water levels on Bay Springs are controlled by Tennessee Valley Authority's operations at Pickwick Dam.

Location: Within Tishomingo and Prentiss Counties approximately 26 miles East of Booneville on Hwy 4.

Fishery Management: Largemouth, Spotted, and Smallmouth Bass, crappie, bream, and catfish.

Purchase a Fishing License: <u>https://www.ms.gov/mdwfp/hunting_fishing/</u>

Amenities

- 7 public boat ramps
- Marina
- Campground
- Public Beach
- Picnic Areas
- Restrooms
- Lock & Dam scenic overlook
- Interpretive center

Regulations

- Black bass (largemouth, spotted, and smallmouth): 10 bass (combined) per day per angler.
- Crappie: 9 inch minimum length limit and 30 crappie per day per angler.
- Bream: 100 bream (combined) per day per angler.
- Catfish: No limit.
- Other creel limits can be found at <u>http://www.mdwfp.com</u>

Fish Stockings

2023 Florida Largemouth Bass (100,000)

2019 Florida Largemouth Bass (110,000)

<u>2013</u> Florida Largemouth Bass (75,000)

<u>2011</u> Florida Largemouth Bass (75,000)

2009 Florida Largemouth Bass (114,660)

> <u>1999</u> Threadfin Shad (60,000)

Fishing Tips

<u>Black Bass</u>

- Target main lake and secondary points with C/T rigged plastics.
- In the spring, target timber areas in the back of coves with spinner baits.

<u>Crappie</u>

- Try standing timber near Jackson Camp.
- Fish Ashcraft Holler and McDougal Branch using roadrunners or hair jigs.
- Target main lake points in the winter.

<u>Bream</u>

• Fish similar areas as crappie using redworms and crickets.

<u>Catfish</u>

- Target cats at night on the shallow flats with cutbait or liver.
- During the summer, fish the rip-rap on the north side of the lake near Crow's Neck.


Below: Results from the 2021 fall electrofishing survey on Bay Springs. Boat electrofishing was completed along 9.3 miles of shoreline in November. Healthy bass (largemouth and spotted) populations were seen during the survey.


Species	# of Fish Sampled	% of Sample	Minimum Length (Inches)	Average Length (inches)	Maximum Length (inches)	Catch Rate Adult fish (fish/mile)
Gizzard Shad	61	6	10.2	12.2	13.8	7
Threadfin Shad	9	<1	2.8	2.9	3.2	-
Chain Pickerel	9	<1	10.6	16.1	23.1	1
Golden Shiner	29	3	2.8	5.7	9.1	-
Channel Catfish	9	<1	15.8	17.8	20.6	1
Bluegill	450	47	0.2	3.7	7.7	35
Longear Sunfish	52	5	1.6	3.5	4.7	4
Redear Sunfish	141	15	1.6	6.3	9.6	13
Smallmouth Bass	4	<1	8.5	13.3	17.4	<1
Spotted Bass	49	5	2.4	9.7	17	4
Largemouth Bass	103	11	4.7	12.8	21.7	9
Crappie (White and Black)	10	1	2.6	5.2	12.1	<1
Yellow Perch	35	4	2.8	6.6	11.1	2


Below: Bay Springs has a healthy crappie population. These Black Crappie were collected from a single piece of woody debris, and almost all were over the 9" minimum length limit.




Below: Length distributions for Largemouth Bass, Spotted Bass, Bluegill, and Redear Sunfish collected during fall 2021 electrofishing. Sampling was conducted over 9.3 miles of shoreline in November.












Right: Percentage of anglers targeting bream, crappie, black bass, and catfish at Bay Springs Lake. The percentages of anglers targeting each species have varied over the last few years; however, crappie anglers have increased in total percentage over the long-term.





Left top: Angler catch rates for bream, crappie, black bass, and catfish at Bay Springs based on spring creel surveys. Bream anglers saw a significant increase in catch rates in 2018. Catch rates for bass and crappie were similar to the historical averages. It is possible that the increase in aquatic vegetation has spread fish out more, making it more difficult for anglers to find concentrations of fish.



Left bottom: The percentage of fish caught that are harvested by anglers at Bay Springs Lake during spring creel surveys.

Have you been interviewed on Bay Springs in 2022?	%	
Yes	3	
No	97	
Do you read and/or plan your fishing trips around the MDWFP weekly fishing reports?	%	
Yes	26	
No	74	
Would you support instituting a 14" minimum length limit on black bass throughout the TTW?	%	
Support	35	
Not Support	12	
No Opinion	53	
Considering all factors, how satisfied are you with the fishing at Bay Springs?		
Very Satisfied	50	
Satisfied	35	
Neutral	12	
Dissatisfied	3	
Very Dissatisfied	0	



Left: Creel surveys provide a good opportunity to collect information and opinions from anglers. Ninety-seven percent of the anglers were interviewed only one time during 2022, which shows that many different anglers fish at Bay Springs. Twenty-six percent said they read the weekly MDWFP fishing reports for Bay Springs. The majority of bass anglers had no opinion on instituting a 14" minimum length limit on black bass throughout the Tenn-Tom Waterway, and eighty-five percent of the anglers interviewed were satisfied with the fishing at Bay Springs Lake.

Right: Average length of crappie and black bass harvested by anglers at Bay Springs Lake during spring creel surveys. Although the harvest rate for bass has decreased (none were kept during 2022 surveys), the size of bass harvested has increased. In contrast, crappie anglers may be harvesting more crappie due to a decrease in the size of crappie caught.

